

On the Eigenvalues of Certain Matrices Over $\mathbb{Z}_m$

Liang Feng Zhang
Nanyang Technological University
liangfeng.zhang@ntu.edu.sg

Abstract

Let $m, n > 1$ be integers and $\mathbb{P}_{n,m}$ be the point set of the projective $(n-1)$ -space (defined by [2]) over the ring $\mathbb{Z}_m$ of integers modulo m . Let $A_{n,m} = (a_{uv})$ be the matrix with rows and columns being labeled by $\mathbb{P}_{n,m}$, where $a_{uv} = 1$ if the inner product $\langle u, v \rangle = 0$ and $a_{uv} = 0$ otherwise. Let $B_{n,m} = A_{n,m} A_{n,m}^t$. The eigenvalues of $B_{n,m}$ have been studied by [1, 2, 3], where their applications in the study of expanders and locally decodable codes were described. In this paper, we completely determine the eigenvalues of $B_{n,m}$ for general integers m and n .

1 Introduction

Let $m > 1$ be an integer. Let $\mathbb{Z}_m$ be the ring of integers modulo m and $\mathbb{Z}_m^*$ be the group of units of $\mathbb{Z}_m$. Let $n > 1$ be an integer and $\mathbb{Z}_m^n$ be the set of n -tuples with entries in $\mathbb{Z}_m$. We say that $u, v \in \mathbb{Z}_m^n$ are equivalent (and write $u \sim v$) if there is a $\lambda \in \mathbb{Z}_m^*$ such that $u_i = \lambda v_i$ for every $i \in [n]$. If u is not equivalent to v , we write $u \not\sim v$. Let $\mathbb{S}_{n,m} = \{u \in \mathbb{Z}_m^n : \gcd(u_1, u_2, \dots, u_n, m) = 1\}$ and

$$\mathbb{P}_{n,m} \triangleq \mathbb{S}_{n,m} / \sim \quad (1)$$

be the set of equivalence classes of elements of $\mathbb{S}_{n,m}$ under $\sim$. Let $A_{n,m} = (a_{uv})$ be the matrix with rows and columns being labeled by $\mathbb{P}_{n,m}$, where $a_{uv} = 1$ if the inner product $\langle u, v \rangle = 0$ and $a_{uv} = 0$ otherwise. Let $B_{n,m} = A_{n,m} A_{n,m}^t$. For every $u \in \mathbb{P}_{n,m}$, let $N(u)$ be the neighborhood of u . Let $\theta_{n,m} = |\mathbb{P}_{n,m}|$. Chee et al. [2] showed that

$$\theta_{n,m} = m^{n-1} \prod_{p|m} \left(1 + \frac{1}{p} + \dots + \frac{1}{p^{n-1}} \right) \quad (2)$$

and $|N(u)| = \theta_{n-1,m}$ for every $u \in \mathbb{P}_{n,m}$.

Let a, b be positive integers. We denote by I_a the identity matrix of order a . We denote by $O_{a \times b}$ and $J_{a \times b}$ the $a \times b$ all-zero and all-one matrices, respectively. In particular, we write O_a and J_a when $a = b$, and write I , O and J when a, b are obvious. When m is a prime, Alon [1] showed that $B_{n,m}$ has two distinct eigenvalues $\theta_{n-1,m}^2$ and m^{n-2} . The eigenvectors with eigenvalue $\theta_{n-1,m}^2$ is the single column of $J_{l \times 1}$ and the eigenvectors with eigenvalue m^{n-2} are the columns of the matrix

$$R_d = \begin{pmatrix} I_d \\ -J_{1 \times d} \end{pmatrix}, \quad (3)$$

where $l = \theta_{n,m}$ and $d = \theta_{n,m} - 1$. When $m = pq$ for two distinct primes, the eigenvalues of $B_{n,m}$ have been determined by Chee et al. [3] (see Lemma 3.2), which have important applications in the study of matching families in $\mathbb{Z}_m^n$. Our work in this paper is mainly motivated by Chee et al. [3] for its potential applications in the study of matching families.

2 Results

In this paper, we completely determine the eigenvalues of $B_{n,m}$ for any positive integer m . First of all, we deal with the prime power case and show the following theorem.

Theorem 2.1 (Prime Power Case) *Let $m = p^e$ for a prime p and positive integers e and n . Then the eigenvalues of $B_{n,m}$ and their multiplicities are as follows:*

<i>Eigenvalue</i>	<i>Multiplicity</i>
$\lambda_1 = p^{2(e-1)(n-2)} \cdot \theta_{n-1,p}^2$	$d_1 = 1$
$\lambda_2 = p^{(2e-1)(n-2)}$	$d_2 = \theta_{n,p} - 1$
$\lambda_s = p^{(2e+1-s)(n-2)}$	$d_s = (p^{n-1} - 1)\theta_{n,p^{s-2}}$

Table 1: The eigenvalues of the matrix $B_{n,m}$

Theorem 3.1 makes it very easy to determine the eigenvalues of $B_{n,m}$ for a general integer m . We define the tensor product of two matrices $A = (a_{ij})$ and B to be the block matrix

$$A \otimes B = (a_{ij} \cdot B). \quad (4)$$

We say that $A \sim B$ if A can be obtained from B by permutating rows and columns. Clearly, if $A \sim B$, then A and B have the same eigenvalues. The following lemma allows us to determine the eigenvalues of $B_{n,m}$ via Theorem 3.1.

Lemma 2.1 (Tensor Lemma) *Let $m = m_1 \cdots m_r = p_1^{e_1} \cdots p_r^{e_r}$ for distinct primes $p_1, \dots, p_r$ and positive integers $e_1, \dots, e_r$, where $m_s = p_s^{e_s}$ for every $s \in [r]$. Then we have that*

$$B_{n,m} \sim B_{n,m_1} \otimes \cdots \otimes B_{n,m_r}. \quad (5)$$

Theorem 2.2 (General Case) *Let $m = m_1 \cdots m_r = p_1^{e_1} \cdots p_r^{e_r}$ for distinct primes $p_1, \dots, p_r$ and positive integers $e_1, \dots, e_r$, where $m_s = p_s^{e_s}$ for every $s \in [r]$. Let λ_s be an eigenvalue of B_{n,m_s} of multiplicity d_s for every $s \in [r]$. Then $\lambda_1 \cdots \lambda_r$ is an eigenvalue of $B_{n,m}$ of multiplicity $d_1 \cdots d_r$.*

3 The Prime Power Case

In this section, we determine the eigenvalues of $B_{n,m}$ whenever m is a prime power.

3.1 Linear Equation Systems Over $\mathbb{Z}_{p^e}$

Let $a, b, c, d \in \mathbb{Z}_{p^e}$ for a prime p and an integer $e > 0$. We consider the following equation system

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} \equiv \begin{pmatrix} 0 \\ 0 \end{pmatrix} \pmod{p^e}. \quad (6)$$

Let $\mathcal{N}$ be the number of pairs $(x, y) \in \mathbb{Z}_{p^e}^2$ that satisfy (6). Let

$$N = \gcd(ad - bc, p^e \cdot \gcd(a, b, c, d, p^e)). \quad (7)$$

Lemma 3.1 *We have that $\mathcal{N} = N$.*

Proof: Suppose that $\gcd(ad - bc, p^e) = p^f$ and $\gcd(a, b, c, d, p^e) = p^g$ for $f, g \in \{0, 1, \dots, e\}$.

- If $g = e$, then $a = b = c = d \equiv 0 \pmod{p^e}$. It follows that $\mathcal{N} = p^{2e} = N$.
- If $g = 0$, then (w.l.o.g.) we may suppose that $\gcd(a, p) = 1$. Let a^{-1} be the inverse of a modulo p^e . Then it is easy to see that $(x, y) \in \mathbb{Z}_{p^e}^2$ satisfies (6) if and only if

$$y \equiv 0 \pmod{p^{e-f}} \text{ and } x \equiv -a^{-1}by \pmod{p^e}.$$

Hence, $\mathcal{N}$ is equal to the number of the choices of $y \in \mathbb{Z}_{p^e}$, which is equal to $p^f = N$.

- If $0 < g < e$, then there are $p^{-2g}N$ pairs $(x, y) \in \mathbb{Z}_{p^{e-g}}^2$ that satisfy

$$p^{-g} \cdot \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} \equiv \begin{pmatrix} 0 \\ 0 \end{pmatrix} \pmod{p^{e-g}} \quad (8)$$

due to the previous case. If $(x, y) \in \mathbb{Z}_{p^e}^2$ is a solution for (6), then

$$(x \bmod p^{e-g}, y \bmod p^{e-g})$$

is a solution for (8). Conversely, if $(x, y) \in \mathbb{Z}_{p^{e-g}}^2$ is a solution for (8), then

$$(x + kp^{e-g}, y + lp^{e-g})$$

is a solution of (6) for every $k, l \in \mathbb{Z}_{p^g}$. Hence, $\mathcal{N} = p^{2g} \cdot (p^{-2g} \cdot N) = N$.

□

Definition 3.1 (*p*-adic Valuation) *Let p be a prime and $t \in \mathbb{Z}$, we denote by $\nu_p(t)$ the largest nonnegative integer e such that $p^e | t$. In particular, we set $\nu_p(0) = \infty$.*

Lemma 3.2 *Let p be a prime and $s, t \in \mathbb{Z}$. Then*

- $\nu_p(t) \leq \infty$ and $\nu_p(t) = \infty$ if and only if $t = 0$;
- $\nu_p(st) = \nu_p(s) + \nu_p(t)$;
- $\nu_p(s + t) \geq \min\{\nu_p(s), \nu_p(t)\}$ and the equality holds when $\nu_p(s) \neq \nu_p(t)$.

Let $u, v \in \mathbb{P}_{n, p^e}$ be arbitrary. For every $i, j \in [n]$, we define

$$\xi_{ij} = u_i v_j - u_j v_i. \quad (9)$$

Let

$$\alpha = \min\{\nu_p(\xi_{ij}) : i, j \in [n]\} \text{ and } \xi = p^{\min(\alpha, e)}. \quad (10)$$

Then ξ is the greatest common divisor of all the integers in $\{\xi_{ij} : i, j \in [n]\}$ and p^e .

Lemma 3.3 *The following properties hold*

- (a) *There are distinct integers $i, j \in [n]$ such that*

$$\gcd(u_i, u_j, v_i, v_j, p^e) = 1 \text{ and } \xi = \gcd(\xi_{ij}, p^e). \quad (11)$$

- (b) $\nu_p(\xi) < e$.

Proof: Since $u \in \mathbb{P}_{n,p^e}$, there is an integer $s \in [n]$ such that $\gcd(u_s, p) = 1$.

(a) If $\alpha = \infty$, then we may take $i = s$ and then (11) follows. From now on, we suppose that $\alpha < \infty$. If (11) does not hold, then we have that $\gcd(u_i, u_j, v_i, v_j, p^e) > 1$ for any $i, j \in [n]$ such that $\nu_p(\xi_{ij}) = \alpha$. Since $\gcd(u_s, p) = 1$, this implies that $p^{\alpha+1} | \xi_{si}$ and $p^{\alpha+1} | \xi_{sj}$. It follows that $p^{\alpha+1} | \xi_{ij}$, which is a contradiction.

(b) If $\nu_p(\xi) = e$, then $\xi_{si} \equiv 0 \pmod{p^e}$ for every $i \in [n]$. It follows that $v_i \equiv u_s^{-1} v_s u_i \pmod{p^e}$, where u_s^{-1} is the inverse of u_s modulo p^e . Note that $v_s \not\equiv 0 \pmod{p}$ since otherwise we will have that $v \notin \mathbb{P}_{n,p^e}$. It follows that $u \sim v$, which is a contradiction because u and v are distinct equivalence classes. $\square$

We shall determine the (u, v) entry of B_{n,p^e} , i.e., b_{uv} in the remaining of this section. Clearly, b_{uv} is the number of $w \in \mathbb{P}_{n,p^e}$ that satisfies the following equation system:

$$\begin{cases} \langle u, w \rangle \equiv 0 \pmod{p^e}, \\ \langle v, w \rangle \equiv 0 \pmod{p^e}. \end{cases} \quad (12)$$

For every $g \in \{0, 1, \dots, e\}$, we define

$$p^g \cdot \mathbb{Z}_{p^e}^n = \{w \in \mathbb{Z}_{p^e}^n : p^g | \gcd(w_1, \dots, w_n)\},$$

Lemma 3.4 *For every $g \in \{0, 1, \dots, e\}$, the equation system (12) has exactly $p^{\beta+(e-g)(n-2)}$ solutions in $p^g \cdot \mathbb{Z}_{p^e}^n$, where $\beta = \min(\nu_p(\xi), e - g)$.*

Proof: We prove for Case I: $g = 0$ and Case II: $0 < g \leq e$, respectively.

- Case I: Due to Lemma 3.3, we may suppose that $\gcd(u_1, u_2, v_1, v_2, p^e) = 1$ and $\xi = \gcd(\xi_{12}, p^e)$. It suffices to show that for every $(w_3, \dots, w_n) \in \mathbb{Z}_{p^e}^{n-2}$, the following equation system

$$\begin{pmatrix} u_1 & u_2 \\ v_1 & v_2 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} \equiv \begin{pmatrix} -\sum_{k=3}^n u_k w_k \\ -\sum_{k=3}^n v_k w_k \end{pmatrix} \pmod{p^e} \quad (13)$$

has exactly ξ solutions in $\mathbb{Z}_{p^e}^2$. Due to Lemma 3.1, the homogenous form of (13) has exactly ξ solutions in $\mathbb{Z}_{p^e}^2$.

Hence, it suffices to show that (13) is solvable. Let $\nu_p(\xi_{12}) = h$. Then there is an integer $\eta \in \mathbb{Z}_{p^e}^*$ such that $\xi_{12} = p^h \eta$. Let η^{-1} be the inverse of η modulo p^e . Then

$$(x, y) = \left(\sum_{k=3}^n \frac{\eta^{-1} \xi_{2k}}{p^h} w_k, \sum_{k=3}^n \frac{\eta^{-1} \xi_{k1}}{p^h} w_k \right)$$

is a solution of (13).

- Case II: Let $w \in p^g \cdot \mathbb{Z}_{p^e}^n$ and $w' = p^{-g} \cdot w$. Then w satisfies (12) if and only if

$$\begin{cases} \langle u, w' \rangle \equiv 0 \pmod{p^{e-g}}, \\ \langle v, w' \rangle \equiv 0 \pmod{p^{e-g}}. \end{cases} \quad (14)$$

Due to Case I, the equation system (14) has exactly $p^{\beta+(e-g)(n-2)}$ solutions $w' \in \mathbb{Z}_{p^{e-g}}^n$. $\square$

Lemma 3.5 *Let $u, v \in \mathbb{P}_{n,m}$ be arbitrary. Then the (u, v) entry of B_{n,p^e} is*

$$b_{uv} = \frac{1}{\phi(p^e)} \left(p^{\nu_p(\xi)+e(n-2)} - p^{\min(\nu_p(\xi), e-1)+(e-1)(n-2)} \right). \quad (15)$$

Proof: Clearly, we have that $\mathbb{Z}_{p^e}^n \setminus p \cdot \mathbb{Z}_{p^e}^n = \mathbb{S}_{n,p^e}$. Let N_{uv} be the number of solutions of (12) in $\mathbb{S}_{n,p^e}$. Due to Lemma 3.4, it is not hard to see that

$$N_{uv} = \begin{cases} p^{\nu_p(\xi)+e(n-2)} - p^{\min(\nu_p(\xi), e-1)+(e-1)(n-2)} & \text{if } u = v, \\ p^{e(n-1)} - p^{(e-1)(n-1)} & \text{if } u \neq v. \end{cases}$$

Note that $\nu_p(\xi) = e$ when $u = v$ and each equivalence class in $\mathbb{P}_{n,p^e}$ contains exactly $\phi(p^e)$ elements of $\mathbb{S}_{n,p^e}$. It follows that $b_{uv} = \phi(p^e)^{-1} N_{uv}$ and the equation (15) follows. $\square$

3.2 The Eigenvalues

We proceed to determine the eigenvalues of B_{n,p^e} . Let $\sigma : \mathbb{S}_{n,p^e} \rightarrow \mathbb{S}_{n,p^{e-1}}$ be the mapping defined by

$$\sigma(u) = (u_1 \bmod p^{e-1}, \dots, u_n \bmod p^{e-1}), \quad (16)$$

where $u \in \mathbb{S}_{n,p^e}$. Let $\tau : \mathbb{S}_{n,p^{e-1}} \rightarrow \mathbb{P}_{n,p^{e-1}}$ to be the mapping such that

$$\tau(u) = \text{the equivalence class of } u. \quad (17)$$




Figure 1: The mappings

Let $\rho = \tau \circ \sigma$. The following lemma shows that the mapping ρ is balanced in the sense that every equivalence class in $\mathbb{P}_{n,p^{e-1}}$ has the same number of preimages in $\mathbb{S}_{n,p^e}$.

Lemma 3.6 *We have that $|\rho^{-1}(v)| = p^n \phi(p^{e-1})$ for every $v \in \mathbb{P}_{n,p^{e-1}}$.*

Proof: Let $w \in \mathbb{S}_{n,p^{e-1}}$ be arbitrary. For every $\gamma = (\gamma_1, \dots, \gamma_n) \in \mathbb{Z}_p^n$, we define

$$w_\gamma = (w_1 + \gamma_1 p^{e-1}, \dots, w_n + \gamma_n p^{e-1}).$$

It is easy to see that $w_\gamma \in \rho^{-1}(w)$ for every $\gamma \in \mathbb{Z}_p^n$ and $w_\gamma \neq w_{\gamma'}$ whenever $\gamma, \gamma' \in \mathbb{Z}_p^n$ are distinct. It follows that $|\sigma^{-1}(w)| \geq p^n$. If $|\sigma^{-1}(w)| > p^n$ for certain choice of $w \in \mathbb{S}_{n,p^{e-1}}$, then we must have that

$$|\mathbb{S}_{n,p^e}| > p^n |\mathbb{S}_{n,p^{e-1}}|.$$

However, due to (2), we have that

$$|\mathbb{S}_{n,p^e}| = \phi(p^e) |\mathbb{P}_{n,p^e}| = p^n \phi(p^{e-1}) |\mathbb{P}_{n,p^{e-1}}| = p^n |\mathbb{S}_{n,p^{e-1}}|.$$

Therefore, we must have that $|\sigma^{-1}(w)| = p^n$. Let $v \in \mathbb{P}_{n,p^{e-1}}$ be arbitrary. Then it is easy to see that $|\tau^{-1}(v)| = \phi(p^{e-1})$. It follows that $|\rho^{-1}(v)| = |\sigma^{-1}(\tau^{-1}(v))| = p^n \phi(p^{e-1})$. $\square$

Let $\psi : \mathbb{S}_{n,p^e} \rightarrow \mathbb{P}_{n,p^e}$ be the mapping defined by

$$\psi(u) = \text{the equivalence class of } u, \quad (18)$$

where $u \in \mathbb{S}_{n,p^e}$. Let $\delta : \mathbb{P}_{n,p^e} \rightarrow \mathbb{P}_{n,p^{e-1}}$ be the mapping defined by

$$\delta(u) = \text{the equivalence class of } (u_1 \bmod p^{e-1}, \dots, u_n \bmod p^{e-1}), \quad (19)$$

where $u \in \mathbb{P}_{n,p^e}$. Then $\delta \circ \psi = \rho$. The following lemma shows that the mapping δ is p^{n-1} to 1.

Lemma 3.7 *We have that $|\delta^{-1}(v)| = p^{n-1}$ for every $v \in \mathbb{P}_{n,p^{e-1}}$.*

Proof: Suppose that $u, v \in \mathbb{S}_{n,p^e}$ and $u \sim v$. Then there is an integer $\lambda \in \mathbb{Z}_{p^e}^*$ such that $u_i \equiv \lambda v_i \bmod p^e$ for every $i \in [n]$. It follows that $u_i \equiv \lambda v_i \bmod p^{e-1}$ for the integer $\lambda \in \mathbb{Z}_{p^{e-1}}^*$. In other words, we have that $\rho(u) = \rho(v)$. Hence, for every $v \in \mathbb{P}_{n,p^{e-1}}$, the set $\rho^{-1}(v)$ is the union of disjoint equivalence classes of the elements of $\mathbb{S}_{n,p^e}$. Due to Lemma 3.6, we have that $|\rho^{-1}(v)| = p^n \phi(p^{e-1})$. Since the elements in $\rho^{-1}(v)$ consists of $\phi(p^e)^{-1} p^n \phi(p^{e-1}) = p^{n-1}$ equivalence classes, we have that

$$|\delta^{-1}(v)| = \text{the number of equivalence classes contained by } \rho^{-1}(v) = p^{n-1},$$

which is the expected result. $\square$

(001)	6	1	1	1	1	1	1	2	1	1	1	1	1	2	1	1	1	1	1	1	1	2	1	1	1	1	1	1	1
(010)	1	6	1	1	1	1	1	1	2	1	1	1	1	1	2	1	1	1	1	1	1	1	1	2	1	1	1	1	1
(011)	1	1	6	1	1	1	1	1	1	2	1	1	1	1	1	2	1	1	1	1	1	1	1	1	2	1	1	1	1
(100)	1	1	1	6	1	1	1	1	1	1	2	1	1	1	1	1	2	1	1	1	1	1	1	1	1	2	1	1	1
(101)	1	1	1	1	6	1	1	1	1	1	2	1	1	1	1	1	2	1	1	1	1	1	1	1	1	2	1	1	1
(110)	1	1	1	1	1	6	1	1	1	1	1	2	1	1	1	1	1	2	1	1	1	1	1	1	1	1	2	1	1
(111)	1	1	1	1	1	1	6	1	1	1	1	1	2	1	1	1	1	1	2	1	1	1	1	1	1	1	1	2	1
(021)	2	1	1	1	1	1	1	6	1	1	1	1	1	1	2	1	1	1	1	1	1	1	2	1	1	1	1	1	1
(012)	1	2	1	1	1	1	1	1	6	1	1	1	1	1	1	2	1	1	1	1	1	1	1	2	1	1	1	1	1
(013)	1	1	2	1	1	1	1	1	1	6	1	1	1	1	1	1	2	1	1	1	1	1	1	1	2	1	1	1	1
(102)	1	1	1	2	1	1	1	1	1	1	6	1	1	1	1	1	2	1	1	1	1	1	1	1	1	2	1	1	1
(103)	1	1	1	1	2	1	1	1	1	1	1	6	1	1	1	1	2	1	1	1	1	1	1	1	1	2	1	1	1
(112)	1	1	1	1	1	2	1	1	1	1	1	1	6	1	1	1	1	2	1	1	1	1	1	1	1	1	2	1	1
(113)	1	1	1	1	1	1	2	1	1	1	1	1	1	6	1	1	1	1	2	1	1	1	1	1	1	1	1	2	1
(201)	2	1	1	1	1	1	1	2	1	1	1	1	1	1	6	1	1	1	1	1	1	2	1	1	1	1	1	1	1
(210)	1	2	1	1	1	1	1	1	2	1	1	1	1	1	1	6	1	1	1	1	1	1	1	2	1	1	1	1	1
(211)	1	1	2	1	1	1	1	1	1	2	1	1	1	1	1	1	6	1	1	1	1	1	1	1	2	1	1	1	1
(120)	1	1	1	2	1	1	1	1	1	1	2	1	1	1	1	1	1	6	1	1	1	1	1	1	1	2	1	1	1
(121)	1	1	1	1	2	1	1	1	1	1	1	2	1	1	1	1	1	1	6	1	1	1	1	1	1	2	1	1	1
(130)	1	1	1	1	1	2	1	1	1	1	1	1	2	1	1	1	1	1	1	6	1	1	1	1	1	1	2	1	1
(131)	1	1	1	1	1	1	2	1	1	1	1	1	1	2	1	1	1	1	1	1	6	1	1	1	1	1	1	2	1
(221)	2	1	1	1	1	1	1	2	1	1	1	1	1	1	2	1	1	1	1	1	1	2	1	1	1	1	1	1	1
(212)	1	2	1	1	1	1	1	1	2	1	1	1	1	1	1	2	1	1	1	1	1	1	1	2	1	1	1	1	1
(213)	1	1	2	1	1	1	1	1	1	2	1	1	1	1	1	1	2	1	1	1	1	1	1	1	2	1	1	1	1
(122)	1	1	1	2	1	1	1	1	1	1	2	1	1	1	1	1	1	2	1	1	1	1	1	1	1	6	1	1	1
(123)	1	1	1	1	2	1	1	1	1	1	1	2	1	1	1	1	1	1	2	1	1	1	1	1	1	1	6	1	1
(132)	1	1	1	1	1	2	1	1	1	1	1	1	2	1	1	1	1	1	1	2	1	1	1	1	1	1	6	1	1
(133)	1	1	1	1	1	1	2	1	1	1	1	1	1	2	1	1	1	1	1	1	2	1	1	1	1	1	1	6	1

Figure 2: The matrix $B_{3,4}$

Due to Lemma 3.7, we can partition $\mathbb{P}_{n,p^e}$ as $l = p^{n-1}$ disjoint subsets $K_1, \dots, K_l$ such that

$$|K_h \cap \delta^{-1}(u)| = 1. \quad (20)$$

for every $h \in [l]$ and $u \in \mathbb{P}_{n,p^{e-1}}$. For every $a, b \in [l]$, we denote by C_{ab} a matrix with rows and columns being labeled by K_a and K_b , respectively. For every $u \in K_a$ and $v \in K_b$, the (u, v) entry of C_{ab} is defined to be the number of solutions of (12) in $\mathbb{P}_{n,p^e}$. Let $C = (C_{ab})$ be a block matrix. Then

$$C \sim B_{n,p^e} \quad (21)$$

Example 3.1 We explain the above description by an example. Let $n = 3$, $p = 2$ and $e = 2$. Then simple calculations show that $\mathbb{P}_{3,4}$ consists of the following equivalence classes

$$\begin{aligned} K_1 : & 001 \quad 010 \quad 011 \quad 100 \quad 101 \quad 110 \quad 111 \\ K_2 : & 021 \quad 012 \quad 013 \quad 102 \quad 103 \quad 112 \quad 113 \\ K_3 : & 201 \quad 210 \quad 211 \quad 120 \quad 121 \quad 130 \quad 131 \\ K_4 : & 221 \quad 212 \quad 213 \quad 122 \quad 123 \quad 132 \quad 133 \end{aligned} \quad (22)$$

where $l = 4$ and the i th row is the i th subset of $\mathbb{P}_{3,4}$ for every $i \in \{1, 2, 3, 4\}$. Clearly, we have that $\mathbb{P}_{3,2} = \{001, 010, 011, 100, 101, 110, 111\}$. It is trivial to verify that $|\delta^{-1}(u) \cap K_h| = 1$ for every $u \in \mathbb{P}_{3,2}$ and $h \in \{1, 2, 3, 4\}$. Figure 2 depicts the matrix $B_{3,4}$, where the rows are columns are labeled by elements in $\mathbb{P}_{3,4}$. Actually, the $B_{3,4}$ is a 4×4 block matrix, where each block is a square matrix of order 7. More precisely, we have that

$$C_{aa} = \begin{pmatrix} 6 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 6 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 6 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 6 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 6 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 & 6 & 1 \\ 1 & 1 & 1 & 1 & 1 & 1 & 6 \end{pmatrix} \quad (23)$$

for every $a \in \{1, 2, 3, 4\}$ and

$$C_{ab} = \begin{pmatrix} 2 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 2 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 2 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 2 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 2 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 & 2 & 1 \\ 1 & 1 & 1 & 1 & 1 & 1 & 2 \end{pmatrix} \quad (24)$$

for every $a, b \in \{1, 2, 3, 4\}$ such that $a \neq b$. It is trivial to verify that the (u, v) entry of C_{aa} and C_{ab} is equal to the number of solutions of (12) in $\mathbb{P}_{3,4}$ for every $u \in K_a$ and $v \in K_b$.

$$\begin{aligned} (001) & \parallel 3 & 1 & 1 & 1 & 1 & 1 \\ (010) & \parallel 1 & 3 & 1 & 1 & 1 & 1 \\ (011) & \parallel 1 & 1 & 3 & 1 & 1 & 1 \\ (100) & \parallel 1 & 1 & 1 & 3 & 1 & 1 \\ (101) & \parallel 1 & 1 & 1 & 1 & 3 & 1 \\ (110) & \parallel 1 & 1 & 1 & 1 & 1 & 3 \\ (111) & \parallel 1 & 1 & 1 & 1 & 1 & 3 \end{aligned}$$

Figure 3: The matrix $B_{3,2}$

Note that the matrix $B_{3,2}$ can be depicted by Figure 3. The following lemma shows that there are connections between B_{n,p^e} and $B_{n,p^{e-1}}$. The connections are clear when we partition the set $\mathbb{P}_{n,p^e}$ as the disjoint subsets $K_1, \dots, K_l$, where $\delta(K_h) = \mathbb{P}_{n,p^{e-1}}$ for every $h \in [l]$.

Lemma 3.8 *Let $a, b \in [l]$ and $a \neq b$. Let $u \in K_a$ and $v \in K_b$ be such that $\delta(u) = \delta(v) = w \in \mathbb{P}_{n,p^{e-1}}$. Then the (u, v) entry of C_{ab} is equal to*

$$\frac{1}{\phi(p^e)} \left(p^{e(n-1)-1} - p^{(e-1)(n-1)} \right). \quad (25)$$

Proof: Without loss of generality, we may suppose that $\gcd(w_1, p) = 1$. Since $\delta(u) = \delta(v) = w$, there are integers $x_2, \dots, x_n, y_2, \dots, y_n \in \mathbb{Z}_p$ such that

$$\begin{aligned} u &\sim (w_1, w_2 + x_2 p^{e-1}, \dots, w_n + x_n p^{e-1}), \\ v &\sim (w_1, w_2 + y_2 p^{e-1}, \dots, w_n + y_n p^{e-1}). \end{aligned} \quad (26)$$

Let ξ be defined by (10). Due to (26), it is trivial to verify that $\xi = p^{e-1}$. It follows that (25) is an immediate consequence of (15). $\square$

Lemma 3.9 *Let $a, b \in [l]$ be arbitrary. Let $u \in K_a$ and $v \in K_b$ be such that $\delta(u) = u' \neq v' = \delta(v)$. Then the (u, v) entry of C_{ab} is equal to the product of p^{n-3} and the (u', v') entry of $B_{n,p^{e-1}}$.*

Proof: Clearly, there are integers $x_1, \dots, x_n, y_1, \dots, y_n \in \mathbb{Z}_p$ such that

$$\begin{aligned} u &\sim (u'_1 + x_1 p^{e-1}, u'_2 + x_2 p^{e-1}, u'_3 + x_3 p^{e-1}, \dots, u'_n + x_n p^{e-1}), \\ v &\sim (v'_1 + y_1 p^{e-1}, v'_2 + y_2 p^{e-1}, v'_3 + y_3 p^{e-1}, \dots, v'_n + y_n p^{e-1}). \end{aligned} \quad (27)$$

Let $\eta_{ij} = u'_i v'_j - u'_j v'_i$ for every $i, j \in [n]$. Let ξ_{ij} be defined by (9). Then $\xi_{ij} \equiv \eta_{ij} \pmod{p^{e-1}}$. Since $u \not\sim v$, Lemma 3.3 implies that $\nu_p(\xi) < e$ and $\nu_p(\eta) < e - 1$. We claim that $\nu_p(\xi) = \nu_p(\eta)$. In fact, by Lemma 3.3, there are integers $\hat{i}, \hat{j} \in [n]$ and $\hat{i}, \hat{j} \in [n]$ such that

$$\xi = \gcd(\xi_{\hat{i}\hat{j}}, p^e) \text{ and } \eta = \gcd(\eta_{\hat{i}\hat{j}}, p^{e-1}).$$

Since $\xi_{\hat{i}\hat{j}} \equiv \eta_{\hat{i}\hat{j}} \pmod{p^{e-1}}$, we must have that $\nu_p(\xi_{\hat{i}\hat{j}}) \leq \nu_p(\eta_{\hat{i}\hat{j}})$. It follows that

$$\nu_p(\xi) = \nu_p(\xi_{\hat{i}\hat{j}}) \leq \nu_p(\eta_{\hat{i}\hat{j}}) \leq \nu_p(\eta) = \nu_p(\eta).$$

Similarly, we must have that $\nu_p(\eta) \leq \nu_p(\xi)$. Hence, our claim holds. In particular, we have that $\nu_p(\xi) = \nu_p(\eta) < e - 1$. Now the lemma is an immediate consequence of (15). $\square$

Lemma 3.10 *Let p be a prime and $n, e > 1$ be integers. Then for any integers $a, b \in [l]$, we have that*

$$C_{ab} = \begin{cases} p^{n-3} B_{n,p^{e-1}} - p^{(e-1)(n-2)-1} I & \text{if } a \neq b, \\ p^{n-3} B_{n,p^{e-1}} + \left(p^{e(n-2)} - p^{(e-1)(n-2)-1} \right) I & \text{if } a = b. \end{cases} \quad (28)$$

Proof: As depicted by Example 3.1, we partition the set $\mathbb{P}_{n,p^e}$ as l disjoint subsets, where $l = p^{n-1}$. Equation (15) shows that the diagonal entries of $B_{n,p^{e-1}}$ are all equal to

$$\frac{1}{\phi(p^{e-1})} \left(p^{(e-1)(n-1)} - p^{(e-2)(n-1)} \right).$$

When $a \neq b$, Lemma 3.8 shows that the diagonal entries of C_{ab} are all equal to

$$\frac{1}{\phi(p^e)} \left(p^{e(n-1)-1} - p^{(e-1)(n-1)} \right).$$

Clearly, the difference between the diagonal entries is $p^{(e-1)(n-2)-1}$. Lemma 3.9 shows that the non-diagonal entries of C_{ab} are p^{n-3} times of those of $B_{n,p^{e-1}}$. It follows that

$$C_{ab} = p^{n-3} B_{n,p^{e-1}} - p^{(e-1)(n-2)-1} I,$$

which is the first equality. Note that the diagonal entries of C_{aa} are equal to

$$\frac{1}{\phi(p^e)} \left(p^{e(n-1)} - p^{(e-1)(n-1)} \right).$$

Similarly, we can prove the second part of (28). □

Lemma 3.11 *If λ is an eigenvalue of $B_{n,p^{e-1}}$, then $p^{2n-4} \cdot \lambda$ is an eigenvalue of B_{n,p^e} .*

Proof: Due to Lemma 3.10, $B_{n,p^e} - p^{e(n-2)} I$ is an $l \times l$ block matrix, where each block is equal to

$$C_{12} = p^{n-3} B_{n,p^{e-1}} - p^{(e-1)(n-2)-1} I.$$

It follows that $\mu = p^{n-3} \lambda - p^{(e-1)(n-2)-1}$ is an eigenvalue of B_{n,p^e} . Thus

$$l\mu + p^{e(n-2)} = p^{n-1} \mu + p^{e(n-2)} = p^{2n-4} \lambda$$

is an eigenvalue of B_{n,p^e} . □

Lemma 3.12 *$\lambda_{e+1} = p^{e(n-2)}$ is an eigenvalue of B_{n,p^e} of multiplicity at least $(p^{n-1} - 1)\theta_{n,p^{e-1}}$.*

Proof: Let $a \in [l-1]$. For every $u \in K_a$ and $v \in K_l$ such that $\delta(u) = \delta(v)$, let w be a vector with coordinates being labeled by $\mathbb{P}_{n,p^e}$ such that the coordinate labeled by u is 1, the coordinate labeled by v is -1 and all the other coordinates are 0. Due to Lemma 3.10, we have that $B_{n,p^e} \cdot w = p^{e(n-2)} \cdot w$. Since there are

$$(l-1) \cdot |K_a| = (p^{n-1} - 1) \cdot \theta_{n,p^{e-1}}$$

choices for w when a is taken over $[l-1]$ and u is taken over K_a for every a . Clearly, all the w 's are linearly independent. The eigenvalue is of multiplicity at least $(p^{n-1} - 1)\theta_{n,p^{e-1}}$. □

Theorem 3.1 (Prime Power Case) *Let $m = p^e$ for a prime p and positive integers e and n . Then the eigenvalues of $B_{n,m}$ and their multiplicities are as follows ($s \in \{3, \dots, e+1\}$):*

<i>Eigenvalue</i>	<i>Multiplicity</i>
$\lambda_1 = p^{2(e-1)(n-2)} \cdot \theta_{n-1,p}^2$	$d_1 = 1$
$\lambda_2 = p^{(2e-1)(n-2)}$	$d_2 = \theta_{n,p} - 1$
$\lambda_s = p^{(2e+1-s)(n-2)}$	$d_s = (p^{n-1} - 1)\theta_{n,p^{s-2}}$

Table 2: The eigenvalues of the matrix $B_{n,m}$

Proof: The matrix $B_{n,p}$ has two eigenvalues $\mu_1 = \theta_{n-1,p}^2$ and $\mu_2 = p^{n-2}$, which are of multiplicity $d_1 = 1$ and $d_2 = \theta_{n,p} - 1$, respectively. Lemma 3.11 shows that

- $\lambda_1 = p^{(2n-4)(e-1)} \mu_1$ is an eigenvalue of B_{n,p^e} of multiplicity at least d_1 ;

– $\lambda_2 = p^{(2n-4)(e-1)}\mu_2$ is an eigenvalue of B_{n,p^e} of multiplicity at least d_2 .

Lemma 3.12 shows that $\mu_s = p^{(s-1)(n-2)}$ is an eigenvalue of $B_{n,p^{s-1}}$ of multiplicity at least $d_s = (p^{n-1} - 1)\theta_{n,p^{s-2}}$ for every $s \in \{3, \dots, e+1\}$. Due to Lemma 3.11,

– $\lambda_s = p^{(2n-4)(e-s+1)}\mu_s$ is an eigenvalue of B_{n,p^e} of multiplicity at least d_s for $s \in \{3, \dots, e+1\}$.

The sum of the multiplicities of $\lambda_1, \dots, \lambda_{e+1}$ is at least

$$\sum_{s=1}^{e+1} d_s = 1 + \theta_{n,p} - 1 + \sum_{s=3}^{e+1} (p^{n-1} - 1) \theta_{n,p^{s-2}} = \theta_{n,p^e}$$

Hence, the multiplicity of λ_s must be $d_s = (p^{n-1} - 1)\theta_{n,p^{s-2}}$ for every $s \in [e+1]$. $\square$

4 The General Case

In this section, we determine the eigenvalues of $B_{n,m}$ for a general integer m . Firstly, we show a tensor lemma on the matrix $B_{n,m}$.

Lemma 4.1 *If $m = m_1 m_2$ for two coprime integers m_1 and m_2 , then we have that*

$$B_{n,m} \sim B_{n,m_1} \otimes B_{n,m_2}. \quad (29)$$

Proof: Let $\pi : \mathbb{P}_{n,m_1} \times \mathbb{P}_{n,m_2} \rightarrow \mathbb{P}_{n,m}$ be the mapping defined by $\pi(u, v) = w$, where

$$w_i \equiv u_i \pmod{m_1} \text{ and } w_i \equiv v_i \pmod{m_2} \quad (30)$$

for every $i \in [n]$. The π is well-defined. In fact, let $w = \pi(u, v)$ and $w' = \pi(u', v')$, where $u, u' \in \mathbb{P}_{n,m_1}$ and $v, v' \in \mathbb{P}_{n,m_2}$. If $u \sim u'$ and $v \sim v'$, then there are integers $\lambda \in \mathbb{Z}_{m_1}^*$ and $\mu \in \mathbb{Z}_{m_2}^*$ such that

$$u'_i \equiv \lambda u_i \pmod{m_1} \text{ and } v'_i \equiv \mu v_i \pmod{m_2}. \quad (31)$$

for every $i \in [n]$. Let δ be an integer such that

$$\delta \equiv \lambda \pmod{m_1} \text{ and } \delta \equiv \mu \pmod{m_2} \quad (32)$$

Due to (30), (31), (32), we have that $w'_i \equiv \delta w_i \pmod{m}$ for every $i \in [n]$. Hence, $w' \sim w$.

It is easy to see that the mapping π is bijective and $\theta_{n,m} = \theta_{n,m_1} \theta_{n,m_2}$. Hence, π is bijective. Let w, w' be defined as above. Clearly, $\langle w, w' \rangle \equiv 0 \pmod{m}$ if and only if $\langle u, u' \rangle \equiv 0 \pmod{m_1}$ and $\langle v, v' \rangle \equiv 0 \pmod{m_2}$. Hence, the (w, w') entry of $A_{n,m}$ is equal to 1 if and only if the (u, u') entry of A_{n,m_1} and the (v, v') entry of A_{n,m_2} are both equal to 1. Hence, $A_{n,m} \sim A_{n,m_1} \otimes A_{n,m_2}$. Thus

$$\begin{aligned} B_{n,m} &= A_{n,m} A_{n,m}^t \\ &\sim (A_{n,m_1} \otimes A_{n,m_2})(A_{n,m_1} \otimes A_{n,m_2})^t \\ &= (A_{n,m_1} \otimes A_{n,m_2})(A_{n,m_1}^t \otimes A_{n,m_2}^t) \\ &= (A_{n,m_1} A_{n,m_1}^t) \otimes (A_{n,m_2} A_{n,m_2}^t) \\ &= B_{n,m_1} B_{n,m_2} \end{aligned}$$

which is the expected result. $\square$

As an immediate corollary of Lemma 4.1, we have

Corollary 4.1 Let $m = m_1 \cdots m_r = p_1^{e_1} \cdots p_r^{e_r}$ for distinct primes $p_1, \dots, p_r$ and positive integers $e_1, \dots, e_r$, where $m_s = p_s^{e_s}$ for every $s \in [r]$. Then $B_{n,m} \sim B_{n,m_1} \otimes \cdots \otimes B_{n,m_r}$.

Lemma 4.2 Let λ be an eigenvalue of B_{n,p^e} of multiplicity d . Then there are d eigenvectors (column vectors) $\mathbf{u}_1, \dots, \mathbf{u}_d$ of B_{n,p^e} with eigenvalue λ such that

$$(\mathbf{u}_1, \dots, \mathbf{u}_d) \sim \begin{pmatrix} I_d \\ * \end{pmatrix}. \quad (33)$$

Proof: If $e = 1$, then the single column of $J_{l \times 1}$ is an eigenvector of $B_{n,p}$ with eigenvalue $\theta_{n-1,p}^2$, where $l = \theta_{n,p}$. Furthermore, there are $d = l - 1$ eigenvectors $\mathbf{u}_1, \dots, \mathbf{u}_d$ with eigenvalue p^{n-1} such that

$$(\mathbf{u}_1, \dots, \mathbf{u}_d) = \begin{pmatrix} I_d \\ -J_{1 \times d} \end{pmatrix}.$$

We give the proof for $e > 1$ by induction. Lemma 3.1 shows that B_{n,p^e} has $e + 1$ eigenvalues. We prove for each eigenvalue. Firstly, due to the proof of Lemma 3.12, we have that

- the eigenvectors of B_{n,p^e} with eigenvalue λ_{e+1} have the form (33).

Secondly, for every $s \in [e]$, let $\mathbf{u}_1, \dots, \mathbf{u}_{d_s}$ be the eigenvectors of $B_{n,p^{e-1}}$ with eigenvalue $\lambda_s \cdot p^{-2(n-2)}$. Due to the induction hypothesis, we have that

$$(\mathbf{u}_1, \dots, \mathbf{u}_{d_s}) = \begin{pmatrix} I_{d_s} \\ * \end{pmatrix}.$$

Let $\mathbf{v}_i = J_{p^{n-1} \times 1} \otimes \mathbf{u}_i$ for every $i \in [d_s]$. Lemma 3.10 shows that $\mathbf{v}_1, \dots, \mathbf{v}_{d_s}$ are eigenvectors of B_{n,p^e} with eigenvalue λ_s . Thus, it is easy to see that

- the eigenvectors of B_{n,p^e} with eigenvalue λ_s have the form (33) for every $s \in [e]$.

Finally, by the induction, we have that (33) holds for B_{n,p^e} . □

Theorem 4.1 (General Case) Let $m = m_1 \cdots m_r = p_1^{e_1} \cdots p_r^{e_r}$ for distinct primes $p_1, \dots, p_r$ and positive integers $e_1, \dots, e_r$, where $m_s = p_s^{e_s}$ for every $s \in [r]$. Let λ_s be an eigenvalue of B_{n,m_s} of multiplicity d_s for every $s \in [r]$. Then $\lambda_1 \cdots \lambda_r$ is an eigenvalue of $B_{n,m}$ of multiplicity $d_1 \cdots d_r$.

Proof: Lemma 4.2 shows there are eigenvectors $\mathbf{u}_{s1}, \dots, \mathbf{u}_{sd_s}$ of B_{n,m_s} with eigenvalue λ_s such that

$$(\mathbf{u}_{s1}, \dots, \mathbf{u}_{sd_s}) = \begin{pmatrix} I_{d_s} \\ * \end{pmatrix}. \quad (34)$$

Due to Lemma 2.1, we have that

$$B_{n,m}(\mathbf{u}_{1f_1} \otimes \cdots \otimes \mathbf{u}_{rf_r}) = (B_{n,m_1} \cdot \mathbf{u}_{1f_1}) \otimes \cdots \otimes (B_{n,m_r} \cdot \mathbf{u}_{rf_r}) = (\lambda_1 \cdots \lambda_r) \cdot (\mathbf{u}_{1f_1} \otimes \cdots \otimes \mathbf{u}_{rf_r}),$$

where $1 \leq f_s \leq d_s$ for every $s \in [r]$. Hence, $(\mathbf{u}_{1f_1} \otimes \cdots \otimes \mathbf{u}_{rf_r})$ is an eigenvector of $B_{n,m}$ with eigenvalue $\lambda_1 \cdots \lambda_r$. Due to (34), it is not hard to see that the following eigenvectors

$$\{(\mathbf{u}_{1f_1} \otimes \cdots \otimes \mathbf{u}_{rf_r}) : 1 \leq f_s \leq d_s \text{ for } s \in [r]\}$$

are linearly independent. Hence, $\lambda_1 \cdots \lambda_r$ is an eigenvalue of $B_{n,m}$ of multiplicity at least $d_1 \cdots d_r$.

Let λ_{sj} be the eigenvalue of B_{n,m_s} of multiplicity d_{sj} for every $s \in [r]$ and $j \in [e_s + 1]$. Then $\lambda_{1j_1} \cdots \lambda_{rj_r}$ is an eigenvalue of $B_{n,m}$ of multiplicity at least $d_{1j_1} \cdots d_{rj_r}$. Theorem 3.1 shows that

$$\sum_{j=1}^{e_s+1} d_{sj} = \theta_{n,m_s}$$

for every $s \in [r]$. It follows that

$$\sum_{j_1 \in [e_1+1], \dots, j_r \in [e_r+1]} d_{1j_1} \cdots d_{rj_r} = \sum_{j_1=1}^{e_1+1} d_{1j_1} \cdot \sum_{j_2=1}^{e_2+1} d_{2j_2} \cdots \sum_{j_r=1}^{e_r+1} d_{rj_r} = \prod_{s=1}^r \theta_{n,m_s} = \theta_{n,m}.$$

Hence, the multiplicity of $\lambda_1 \cdots \lambda_r$ cannot be greater than $d_1 \cdots d_r$. In other words, the multiplicity of $\lambda_1 \cdots \lambda_r$ must be exactly $d_1 \cdots d_r$. $\square$

References

- [1] N. Alon. Eigenvalues, geometric expanders, sorting in rounds, and Ramsey theory. *Combinatorica*, 6: 207-219, 1986.
- [2] Y. M. Chee and S. Ling. Projective covering designs. *Bulletin of the London Mathematical Society*, 25(3): 231-239, 1993.
- [3] Y. M. Chee, S. Ling, H. Wang and L. F. Zhang. On the Largest Size of Matching Families in $\mathbb{Z}_{pq}^n$. Manuscript, 2012.